



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>5</sup> :</b> <b>A61K 49/02, 47/12</b>	<b>A1</b>	<b>(11) International Publication Number:</b> <b>WO 93/04702</b> <b>(43) International Publication Date:</b> 18 March 1993 (18.03.93)
<b>(21) International Application Number:</b> PCT/US92/06429 <b>(22) International Filing Date:</b> 31 July 1992 (31.07.92)  <b>(30) Priority data:</b> 751,825                      29 August 1991 (29.08.91)                      US  <b>(71) Applicant:</b> MALLINCKRODT MEDICAL, INC. [US/US]; 675 McDonnell Blvd., P.O. Box 5840, St. Louis, MO 63134 (US).  <b>(72) Inventors:</b> DEUTSCH, Edward, A. ; 12805 Maryland Estates Court, Maryland Heights, MO 63043 (US). GODEMANS, Wilhelmus, Theodorus ; Oude Bergerweg 31, NL-1862 KJ Bergen (NL). DE JONG, Martinus, Theodorus, Maria ; Vangershof 17, NL-1616 PH Hoogkar-spel (NL). MILLER, Kathleen, M. ; 13002 Montmartre Drive, St. Louis, MO 63141 (US). BRODACK, James, W. ; 3827 Kentucky Derby Drive, Florissant, MO 63034 (US).		<b>(74) Agents:</b> KLOSTERMANN, Roy, J. et al.; Mallinckrodt Medical, Inc., 675 McDonnell Blvd., P.O. Box 5840, St. Louis, MO 63134 (US).  <b>(81) Designated States:</b> AU, CA, JP, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, MC, NL, SE).  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
<b>(54) Title:</b> USE OF GENTISIC ACID OR GENTISYL ALCOHOL FOR STABILISING RADIOLABELED PEPTIDES AND PROTEINS		
<div style="display: flex; justify-content: space-around; align-items: center;"><div style="text-align: center;"><chem>OC(=O)c1cc(O)ccc1O</chem> </div><div style="text-align: center;"><chem>OCc1cc(O)ccc1O</chem> </div></div>		
<b>(57) Abstract</b>  Gentisic acid and its derivatives substantially inhibit peptide autoradiolysis. Gentisic acid or its derivatives may also be used in combination with other stabilizers such as inositol, ascorbic acid, and citrate to inhibit autoradiolysis of radiolabeled peptides.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	MN	Mongolia
AU	Australia	FR	France	MR	Mauritania
BB	Barbados	GA	Gabon	MW	Malawi
BE	Belgium	GB	United Kingdom	NL	Netherlands
BF	Burkina Faso	GN	Guinea	NO	Norway
BG	Bulgaria	GR	Greece	NZ	New Zealand
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IE	Ireland	PT	Portugal
CA	Canada	IT	Italy	RO	Romania
CF	Central African Republic	JP	Japan	RU	Russian Federation
CG	Congo	KP	Democratic People's Republic of Korea	SD	Sudan
CH	Switzerland	KR	Republic of Korea	SE	Sweden
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovak Republic
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CS	Czechoslovakia	LU	Luxembourg	SU	Soviet Union
CZ	Czech Republic	MC	Monaco	TD	Chad
DE	Germany	MG	Madagascar	TG	Togo
DK	Denmark	ML	Mali	UA	Ukraine
ES	Spain			US	United States of America

USE OF GENTISIC ACID OR GENTISYL ALCOHOL FOR STABILISING  
RADIOLABELED PEPTIDES AND PROTEINS

BACKGROUND OF THE INVENTION

5        This invention relates to stabilizers for  
radiopharmaceutical compositions. More particularly,  
stabilizers such as gentisic acid and its derivatives,  
alone or in combination with other stabilizers, are used to  
inhibit autoradiolysis of radiolabeled peptides and  
10    proteins.

      The number of therapeutic and diagnostic uses of  
radiolabeled compositions is continually growing. Such  
uses generally involve the introduction of a suitable  
radiolabeled composition into a biological subject.  
15    Detection and imaging of radioactive emissions may be used  
to diagnose and locate the presence of aberrations,  
pathological conditions, and the like. In some cases, the  
radiolabeled composition may be designed to locate in or to  
seek out specific tissues or biological receptors for the  
20    purpose of delivering therapeutic radioactive emissions.

      In general, a radiolabeled composition comprises a  
radionuclide, a carrier agent designed to target the  
specific organ of interest, various auxiliary agents which  
affix the radionuclide to the carrier, a delivery vehicle,  
25    such as water, suitable for injection into or aspiration by  
the patient, physiologic buffers and salts, and the like.

      Some radiopharmaceutical preparations are known to  
require stabilizers. For example, technetium-99m and  
rhenium-186 compositions are unstable in oxygen and require  
30    stabilizers, such as antioxidants or reducing agents, to  
maintain the technetium or rhenium in a usable oxidation  
state. Typical reducing agents used in technetium-99m and  
rhenium-186 compositions include stannous, ferrous, and  
chromous salts. Sometimes other additives, such as  
35    ascorbic acid, d-ascorbic acid, gentisic acid, reductic  
acid, erythorbic acid, p-aminobenzoic acid, 4-  
hydroxybenzoic acid, nicotinic acid, nicotinamide, and 2,5-

dihydroxy-1,4-benzenedisulfonic acid, are included to inhibit the oxidation of the radionuclide or the reducing agent.

Other radionuclides, such as  $^{111}\text{In}$ ,  $^{90}\text{Y}$ , and  $^{67}\text{Ga}$  exist in a stable oxidation state, and therefore, do not require stabilizers to maintain their useful oxidation state.

Over the years, there has been growing interest in preparing radiolabeled proteins such as hormones, macroaggregated albumin ("MAA"), human serum albumin ("HSA"), monoclonal antibodies, or monoclonal antibody fragments for the purpose of diagnosing and treating diseases, such as inflammation, deep vein thrombosis, or cancer. In some cases, autoradiolysis of the labeled protein has been observed. To inhibit or prevent autoradiolysis, experts have suggested adding HSA to the composition (e.g., R.A.J. Kishore, et al., "Autoradiolysis of Iodinated Monoclonal Antibody Preparations," Int. J. Radiat. Appl. Instrum., Part B, Vol. 13, No. 4, pp. 457-459 (1986)) or keeping the radiopharmaceutical composition frozen between preparation and administration (e.g., R.L. Wahl, et al., "Inhibition of Autoradiolysis of Radiolabeled Monoclonal Antibodies by Cryopreservation," J. Nuc. Med., Vol. 31, No. 1, pp. 84-89 (1990)). These techniques for preventing autoradiolysis are often not effective or practical when used with many radiolabeled peptides and proteins.

Recently, a number of exciting new peptides for diagnostic and therapeutic applications have been isolated and synthetically developed. One such peptide is an octapeptide somatostatin analog known as octreotide and described in U.S. Patent No. 4,395,403. Octreotide has a very high binding affinity to somatostatin receptors in a variety of human tumors. By linking octreotide to a suitable chelating agent capable of forming a complex with radionuclides, it has been possible to create radiolabeled

octreotide which effectively images tumors having somatostatin receptors. Somatostatin analogs containing chelating groups are described in UK Patent Publication No. 2,225,579.

5        Despite the potential usefulness of radiolabeled peptides, it has been found that they are very susceptible to autoradiolysis. As used herein, the term autoradiolysis includes chemical decomposition of the peptide by the action of radiation emitting from the radioisotope coupled  
10        to the peptide. Some believe autoradiolysis may be caused by the formation of free radicals, such as hydroxyl radicals, in the water or delivery vehicle by the radiation emitted from the radioisotope.

         From the foregoing, it will be appreciated that what  
15        is needed in the art are stable radiolabeled peptide and protein compositions. Thus, it would be a significant advancement in the art to provide stabilizing agents which substantially inhibit autoradiolysis of radiolabeled peptides and proteins.

20        Such compositions for substantially inhibiting peptide and protein autoradiolysis are disclosed and claimed herein.

#### BRIEF SUMMARY OF THE INVENTION

25        The present invention provides compositions for preparing stable radiolabeled peptide and protein preparations. The stabilizers used in the present invention are able to substantially inhibit autoradiolysis of radiolabeled peptides and proteins. It has been found  
30        that stabilizers that are effective at preventing oxidation of radioisotopes, such as <sup>99m</sup>Tc, are not necessarily effective at preventing autoradiolysis of peptides and proteins. Likewise, it has been found that stabilizers and techniques used to prevent autoradiolysis of proteins, such  
35        as addition of HSA or freezing, are not effective or

practical in many cases. Accordingly, the present invention is directed to compositions containing stabilizers that substantially inhibit autoradiolysis of peptides and proteins.

5       Gentisic acid and its derivatives have been found to be very effective at inhibiting peptide and protein autoradiolysis. Gentisic acid or its derivatives may also be used in combination with other stabilizers, such as inositol and ascorbic acid, to inhibit autoradiolysis.

10

#### DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to compositions containing stabilizers that substantially inhibit autoradiolysis of peptides and proteins. One class of  
15       stabilizers that is very effective at inhibiting autoradiolysis of peptides and proteins is gentisic acid and its derivatives. The radiolabeled peptides and proteins for which the stabilizers of the present invention are needed include peptides and proteins having diagnostic  
20       and therapeutic applications.

Gentisic acid (chemically: 2,5-dihydroxybenzoic acid) is commercially available and can be prepared by several methods known in the art. Derivatives of gentisic acid include the pharmaceutically-acceptable salts and esters of  
25       gentisic acid and gentisyl alcohol. Examples of suitable gentisic acid derivatives are described in U.S. Patent Nos. 4,497,744 and 4,232,000 which are incorporated herein by reference.

Other stabilizers such as inositol and ascorbic acid  
30       may be used in combination with gentisic acid or its derivatives to inhibit autoradiolysis of radiolabeled peptides.

The pharmaceutically-acceptable salts and esters of gentisic acid and gentisic alcohol can be prepared by  
35       standard neutralization and esterification procedures known

in the art, such as the techniques described in U.S. Patent Nos. 4,497,744 and 4,232,000. In practice, the salts and esters of gentisic acid suitable for use in the present invention can be selected according to their solubility.

- 5 Soluble gentisate salts include the soluble alkali metal, alkaline earth metal, heavy metal, and ammonium salts. The alkali metal salts, such as sodium, lithium, and potassium, are very soluble and are currently preferred. The alkaline earth metal gentisate salts, such as calcium and magnesium,  
10 are less soluble, but are still suitable for use herein.

The present invention may be used with a wide range of radioisotopes capable of causing autoradiolysis of peptides. Such radioisotopes include  $\gamma$ -,  $\beta$ -, and  $\alpha$ -emitters.

- 15 Suitable  $\gamma$ -emitters include those radionuclides which are useful for diagnostic techniques. Examples of some typical  $\gamma$ -emitting radionuclides are  $^{67}\text{Ga}$ ,  $^{111}\text{In}$ ,  $^{99\text{m}}\text{Tc}$ ,  $^{169}\text{Yb}$ , and  $^{125}\text{I}$ ,  $^{123}\text{I}$ , and  $^{201}\text{Tl}$ . Examples of possible  $\beta$ -emitting radionuclides include those which are useful in therapeutic  
20 applications such as  $^{90}\text{Y}$ ,  $^{67}\text{Cu}$ ,  $^{186}\text{Re}$ ,  $^{188}\text{Re}$ ,  $^{169}\text{Er}$ ,  $^{121}\text{Sn}$ ,  $^{127}\text{Te}$ ,  $^{143}\text{Pr}$ ,  $^{198}\text{Au}$ ,  $^{109}\text{Pd}$ ,  $^{165}\text{Dy}$ ,  $^{32}\text{P}$ ,  $^{142}\text{Pr}$ ,  $^{177}\text{Lu}$ ,  $^{166}\text{Ho}$ ,  $^{153}\text{Sm}$ ,  $^{90}\text{Y}$ ,  $^{131}\text{I}$ ,  $^{89}\text{Sr}$ , and  $^{105}\text{Rh}$ . Typical  $\alpha$ -emitters include  $^{212}\text{Bi}$ ,  $^{211}\text{At}$ ,  $^{241}\text{Am}$ , and  $^{255}\text{Fm}$ .

- Radiolabeling of peptides and proteins can be achieved  
25 using various methods known in the art. For example, peptides can be labeled through use of a bifunctional chelate, direct labeling, or covalent binding to a specific functional group of an amino acid side chain. The use of a bifunctional chelate involves covalent attachment of a  
30 chelate, which complexes with the radionuclide, to the peptide or protein. Possible bifunctional chelates include DTPA and  $\text{N}_3\text{S}$  ligands. The DTPA may be attached to the peptide or protein by the dicyclic dianhydride method described in U.S. Patent No. 4,479,930, which is  
35 incorporated herein by reference.  $\text{N}_3\text{S}$  ligands may be

attached to the peptide or protein by the methods described in U.S. Patent No. 4,965,392 and European Patent Publication Number 0284071, which are incorporated herein by reference.

5 In direct labeling, the radionuclide binds to the functional group of amino acid side chains present in the peptide or protein. The radionuclide may also bind to reduced forms of a peptide or protein, such as a peptide or protein containing a reduced disulfide bond. One example  
10 of direct labeling known in the art is described in U.S. Patent No. 4,877,868, which is incorporated by reference.

Another well known technique for labeling peptides and proteins involves covalently binding the radionuclide to one specific functional group of an amino acid side chain,  
15 such as incorporation of iodide into the phenol group of a tyrosine residue.

Commercial products for preparing radiopharmaceuticals are generally provided as lyophilized (freeze-dried) "kits" or as liquid formulations. Lyophilized kits are well known  
20 in the art. According to the present invention, lyophilized kits may contain a transfer ligand, such as citric acid, acetic acid, or sodium tartrate, a reducing agent, depending on the radioisotope that is used, a bulking agent, such as inositol or lactose, the peptide or  
25 protein to be labeled, and one or more gentisic acid stabilizers. Additional stabilizers may also be incorporated into the formulation as described herein. The radioisotope is added to the lyophilized kit just prior to patient administration.

30 Liquid formulations usually contain the peptide or protein labeled with the radioisotope. According to the present invention, the liquid formulation also contains one or more gentisic acid derivatives to stabilize the formulation. Other stabilizers, such as inositol and  
35 ascorbic acid, may also be included in the formulation to



improve stability. A surfactant, such as polysorbate 80, and a salt solution to give a desired ionic strength, may also be added to improve stability if the solution is autoclaved. In some cases, the solution may need to be  
5 deoxygenated.

The radiolabeled compositions of the invention may be administered parenterally, preferably intravenously, in the form of injectable pharmaceutical solutions or suspensions according to conventional techniques.

10 Dosages employed in practicing the therapeutic method of the present invention will of course vary depending on the particular condition to be treated, for example the volume of the tumor, the particular chelate employed, the half-life of the radioisotope, and the therapy desired. In  
15 general, the dose is calculated on the basis of radioactivity distribution to each organ and on observed target uptake.

The following examples are offered to further illustrate different aspects of the present invention.  
20 These examples are intended to be purely exemplary and should not be viewed as a limitation on any claimed embodiment.

#### Example 1

##### 25 Preparation of <sup>111</sup>In labeled DTPA-octreotide with citrate additive.

To a lyophilized kit containing 10 µg DTPA-octreotide (N-[3,6,9,9-tetrakis(carboxymethyl)-3,6,9-triazanonanoyl]-D-phenylalanyl-L-hemicystyl-L-phenylalanyl-D-tryptophyl-L-  
30 lysyl-L-threonyl-L-hemicystyl-L-threoninol cyclic (2-7) disulfide, MW=1394.60 gm/mol), 5.6 mg trisodium citrate dihydrate, and 0.4 mg citric acid monohydrate was added 1.0 ml of 0.02 M HCl containing 4.70 mCi of <sup>111</sup>In. The pH of the solution was 4.5. The solution was kept at room  
35 temperature and monitored for the amount of <sup>111</sup>In DTPA-octreotide using reverse-phase HPLC and a Beckman 170

radiometric detector. The HPLC method used a Hamilton PRP-1 column, 25 cm x 4.1 mm, 10 microns and a gradient system, linearly ramping from 100% A (A=10:90 ethanol:water, 10 mM tetrabutyl ammonium phosphate, pH 3) to 43:57 A:B (B=50:50 ethanol:water, 10 mM tetrabutyl ammonium phosphate, pH 3) over 10 minutes, then held at 43:57 A:B for 15 minutes, then linearly ramped to 100% B over 10 minutes. The flow rate was 1.5 ml/min. The retention time of <sup>111</sup>In DTPA-octreotide was 30 minutes.

10 The purity of the <sup>111</sup>In labeled peptide was 87% immediately post-reconstitution and 50% at 15 hours post-reconstitution.

#### Example 2

15 Preparation of <sup>111</sup>In labeled DTPA-octreotide with citrate and inositol additives.

To a lyophilized kit containing 10 µg DTPA-octreotide (N-[3,6,9,9-tetrakis(carboxymethyl)-3,6,9-triazanonanoyl]-D-phenylalanyl-L-hemicystyl-L-phenylalanyl-D-tryptophyl-L-lysyl-L-threonyl-L-hemicystyl-L-threoninol cyclic (2-7) disulfide, MW=1394.60 gm/mol), 5.6 mg trisodium citrate dihydrate, 0.4 mg citric acid monohydrate, and 1.0 mg inositol was added 1.0 ml of 0.02 M HCl containing 6.04 mCi of <sup>111</sup>In. The pH of the solution was 4.5. The solution was kept at room temperature and monitored for the amount of <sup>111</sup>In DTPA-octreotide using reverse-phase HPLC and a Beckman 170 radiometric detector. The HPLC method used a Hamilton PRP-1 column, 25 cm x 4.1 mm, 10 microns and a gradient system, linearly ramping from 100% A (A=10:90 ethanol:water, 10 mM tetrabutyl ammonium phosphate, pH 3) to 100% B (B=50:50 ethanol:water, 10 mM tetrabutyl ammonium phosphate, pH 3) over 60 minutes. The flow rate was 1.5 ml/min. The retention time of <sup>111</sup>In DTPA-octreotide was 44-45 minutes.

The purity of the  $^{111}\text{In}$  labeled peptide was 94% immediately post-reconstitution and 73.5% at 24 hours post-reconstitution.

5

### Example 3

#### Preparation of $^{111}\text{In}$ labeled DTPA-octreotide with citrate, inositol, and ascorbic acid additives.

To a lyophilized kit containing 10  $\mu\text{g}$  DTPA-octreotide (N-[3,6,9,9-tetrakis(carboxymethyl)-3,6,9-triazanonanoyl]-  
10 D-phenylalanyl-L-hemicystyl-L-phenylalanyl-D-tryptophyl-L-lysyl-L-threonyl-L-hemicystyl-L-threoninol cyclic (2-7) disulfide, MW=1394.60 gm/mol), 5.6 mg trisodium citrate dihydrate, and 0.4 mg citric acid monohydrate was added 1.0  
15 mg inositol, 8.8 mg ascorbic acid, 9.9 mg sodium ascorbate, and 1.0 ml of 0.02 M HCl containing 5.05 mCi of  $^{111}\text{In}$ . The pH of the solution was 4.0. The solution was kept at room temperature and monitored for the amount of  $^{111}\text{In}$  DTPA-octreotide using reverse-phase HPLC and a Beckman 170  
20 radiometric detector. The HPLC method used a Hamilton PRP-1 column, 25 cm x 4.1 mm, 10 microns and a gradient system, linearly ramping from 100% A (A=10:90 ethanol:water, 10 mM tetrabutyl ammonium phosphate, pH 3) to 100% B (B=50:50 ethanol:water, 10 mM tetrabutyl ammonium phosphate, pH 3) over 35 minutes. The flow rate was 1.5  
25 ml/min. The retention time of  $^{111}\text{In}$  DTPA-octreotide was 30 minutes.

The purity of the  $^{111}\text{In}$  labeled peptide was 94% immediately post-reconstitution and 86% at 22 hours post-reconstitution.

30

### Example 4

#### Preparation of $^{111}\text{In}$ labeled DTPA-octreotide with citrate, inositol, and gentisic acid additives.

To a lyophilized kit containing 10  $\mu\text{g}$  DTPA-octreotide  
35 (N-[3,6,9,9-tetrakis(carboxymethyl)-3,6,9-triazanonanoyl]-D-phenylalanyl-L-hemicystyl-L-phenylalanyl-D-tryptophyl-L-

lysyl-L-threonyl-L-hemicystyl-L-threoninol cyclic (2-7) disulfide, MW=1394.60 gm/mol), 5.6 mg trisodium citrate dihydrate, and 0.4 mg citric acid monohydrate was added 1.0 mg inositol, 1.5 mg gentisic acid, and 1.0 ml of 0.02 M HCl containing 5.40 mCi of <sup>111</sup>In. The pH of the solution was 4.0. The solution was kept at room temperature and monitored for the amount of <sup>111</sup>In DTPA-octreotide using reverse-phase HPLC and a Beckman 170 radiometric detector. The HPLC method used a Hamilton PRP-1 column, 25 cm x 4.1 mm, 10 microns and a gradient system, linearly ramping from 100% A (A=10:90 ethanol:water, 10 mM tetrabutyl ammonium phosphate, pH 3) to 100% B (B=50:50 ethanol:water, 10 mM tetrabutyl ammonium phosphate, pH 3) over 60 minutes. The flow rate was 1.5 ml/min. The retention time of <sup>111</sup>In DTPA-octreotide was 44-45 minutes.

The purity of the <sup>111</sup>In labeled peptide was 94% immediately post-reconstitution and 94% at 48 hours post-reconstitution.

#### Example 5

##### Preparation of <sup>111</sup>In labeled DTPA-octreotide with gentisic acid additive.

To a vial containing 10 µg DTPA-octreotide (N-[3,6,9,9-tetrakis(carboxymethyl)-3,6,9-triazanonanoyl]-D-phenylalanyl-L-hemicystyl-L-phenylalanyl-D-tryptophyl-L-lysyl-L-threonyl-L-hemicystyl-L-threoninol cyclic (2-7) disulfide, MW=1394.60 gm/mol) in 10 µl of water was added 1 ml of a degassed stock solution of 2.3 mg gentisic acid and 41.4 mg sodium gentisate dihydrate and 25 µl of 0.02 M HCl containing 4.66 mCi of <sup>111</sup>In. The pH of the solution was 4.2. The solution was kept at room temperature and monitored for the amount of <sup>111</sup>In DTPA-octreotide using reverse-phase HPLC and a Beckman 170 radiometric detector. The HPLC method used a Hamilton PRP-1 column, 25 cm x 4.1 mm, 10 microns, eluting with mobile phase A (A=20:10:70

acetonitrile:ethanol:water, 10 mM tetrabutyl ammonium phosphate, pH 3) for 9 minutes, then eluting with mobile phase B (B=50:10:40 acetonitrile:ethanol:water, 10 mM tetrabutyl ammonium phosphate, pH 3) from 9 to 15 minutes.

- 5 The flow rate was 1.2 ml/min. The retention time of <sup>111</sup>In DTPA-octreotide was 9-10 minutes.

The purity of the <sup>111</sup>In labeled peptide was 97% immediately post-reconstitution and 97% at 72 hours post-reconstitution.

10

#### Example 6

##### Preparation of <sup>111</sup>In labeled DTPA-octreotide with acetate and nicotinic acid additives.

- To a lyophilized kit containing 10 µg DTPA-octreotide
- 15 (N-[3,6,9,9-tetrakis(carboxymethyl)-3,6,9-triazanonanoyl]-D-phenylalanyl-L-hemicystyl-L-phenylalanyl-D-tryptophyl-L-lysyl-L-threonyl-L-hemicystyl-L-threoninol cyclic (2→7) disulfide, MW=1394.60 gm/mol) and acetate buffer, pH 4, was added 12.3 mg nicotinic acid and 1.0 ml of 0.02 M HCl
- 20 containing 2.45 mCi of <sup>111</sup>In. The pH of the solution was 3.8. The solution was kept at room temperature and monitored for the amount of <sup>111</sup>In DTPA-octreotide using reverse-phase HPLC and a Beckman 170 radiometric detector. The HPLC method used a Hamilton PRP-1 column, 25 cm x 4.1
- 25 mm, 10 microns and a gradient system, linearly ramping from 100% A (A=10:90 ethanol:water, 10 mM tetrabutyl ammonium phosphate, pH 3) to 100% B (B=50:50 ethanol:water, 10 mM tetrabutyl ammonium phosphate, pH 3) over 60 minutes. The flow rate was 1.5 ml/min. The retention time of <sup>111</sup>In DTPA-
- 30 octreotide was 44-45 minutes.

The purity of the <sup>111</sup>In labeled peptide was 90% immediately post-reconstitution and 80% at 24 hours post-reconstitution.

## Example 7

Preparation of <sup>111</sup>In labeled DTPA-octreotide  
with citrate, inositol, and resorcinol additives.

To a lyophilized kit containing 10 µg DTPA-octreotide  
5 (N-[3,6,9,9-tetrakis(carboxymethyl)-3,6,9-triazanonanoyl]-  
D-phenylalanyl-L-hemicystyl-L-phenylalanyl-D-tryptophyl-L-  
lysyl-L-threonyl-L-hemicystyl-L-threoninol cyclic (2→7)  
disulfide, MW=1394.60 gm/mol), 5.6 mg trisodium citrate  
dihydrate, and 0.4 mg citric acid monohydrate was added 1.0  
10 mg inositol, 11 mg resorcinol, and 1.0 ml of 0.02 M HCl  
containing 5.17 mCi of <sup>111</sup>In. The pH of the solution was  
4.5. The solution was kept at room temperature and  
monitored for the amount of <sup>111</sup>In DTPA-octreotide using  
reverse-phase HPLC and a Beckman 170 radiometric detector.  
15 The HPLC method used a Hamilton PRP-1 column, 25 cm x 4.1  
mm, 10 microns and a gradient system, linearly ramping from  
100% A (A=10:90 ethanol:water, 10 mM tetrabutyl ammonium  
phosphate, pH 3) to 43:57 A:B (B=50:50 ethanol:water, 10 mM  
tetrabutyl ammonium phosphate, pH 3) over 10 minutes, then  
20 held at 43:57 A:B for 15 minutes, then linearly ramped to  
100% B over 10 minutes. The flow rate was 1.5 ml/min. The  
retention time of <sup>111</sup>In DTPA-octreotide was 30 minutes.

The purity of the <sup>111</sup>In labeled peptide was 81%  
immediately post-reconstitution (also, .6% of an <sup>111</sup>In  
25 transfer ligand complex) and 87% at 120 hours post-  
reconstitution.

The results of Examples 1-7 are summarized in Table 1.  
These results demonstrate that gentisic acid, either alone  
or in combination with other stabilizers, is very effective  
30 at preventing radiolysis of radiolabeled peptides.

13

Table 1

	Example	Additive(s)	Initial Purity	Stability	
5	1	citrate	87%	50%	@ 15 hrs
	2	citrate inositol	94%	73.5%	@ 24 hrs
10	3	citrate inositol ascorbic acid	94%	86%	@ 22 hrs
15	4	citrate inositol gentisic acid	94%	94%	@ 48 hrs
	5	gentisic acid	97%	97%	@ 72 hrs
20	6	acetate nicotinic acid	90%	80%	@ 24 hrs
	7	inositol resorcinol	87%	87%	@ 120 hrs

\*81% <sup>111</sup>In DTPA-octreotide and 6% transfer ligand complex.

**Example 8****30      Preparation of <sup>123</sup>I labeled LH-RH.**

This preparation is done in a well-ventilated fume hood equipped with a charcoal filter to absorb any volatile iodine.

In a 1.5 ml polyethylene tube, add 70 μl of buffer A (A=150 mM sodium phosphate, pH 7.4), 10 μl of a 0.1 mM solution of LH-RH (des-gly<sup>10</sup>, [D-ala<sup>6</sup>]-LH-RH ethylamide) in 0.1 M acetic acid, and 5 mCi of <sup>123</sup>I in 5 μl of 0.01 M NaOH. The reaction is started by adding 10 μl of a chloramine T solution (0.5 mg chloramine T/ml buffer A) to the peptide solution and is mixed by cautious aspiration and expulsion of the reaction mixture with the pipet used to add chloramine T. After 1 minute, 100 μl of sodium metabisulfite solution (1 mg sodium metabisulfite/ml buffer A) is added to terminate the reaction and the reaction is

mixed with the pipet as explained previously. The reaction is kept in the hood a few minutes to allow any volatile iodine to exhaust from the reaction vial into the hood.

The solution is loaded onto a PD-10 column (G-25M Sephadex PD-10 column; Pharmacia LKB Biotechnology Inc.; Piscataway, NJ) eluting with PBS, pH 7.4, and 0.5 ml fractions are collected. The  $^{123}\text{I}$  labeled LH-RH is in fractions 6 and 7, which are subsequently combined. Half of this solution is transferred to another vial containing 2 mg of gentisic acid.

The  $^{123}\text{I}$  LH-RH solutions are kept at room temperature and monitored for the amount of iodinated peptide and the amount of free iodide using reverse-phase HPLC and a Beckman 170 radiometric detector. The HPLC method uses a Hamilton PRP-3 column, 15 cm x 4.1 mm, 10 microns and a gradient system eluting from 100% A (A=15:85 acetonitrile:water, 6 mM HCl) to 100% B (B=50:50 acetonitrile:water, 6 mM HCl) over 30 minutes. Free iodide elutes at the void volume in this system while the  $^{123}\text{I}$  LH-RH is retained and has a longer retention time. The solution containing the radiolabeled peptide without the gentisic acid stabilizer shows evidence of autoradiolytic degradation by the appearance of an increasing amount of free iodide in solution, as determined by HPLC analysis.

25

#### Example 9

##### Preparation of $^{186}\text{Re}$ labeled N<sub>3</sub>S-NR-Lu-10.

The NR-Lu-10 monoclonal antibody (NeoRx Corporation, seattle, WA) is labeled with  $^{186}\text{Re}$  using the pre-formed chelate approach developed by NeoRx and described in U.S. Patent No. 4,965,392 and European Patent Publication Number 0284071. The ligand used to form the  $^{186}\text{Re}$  complex is a tetrafluorophenyl ester of [N-(S-ethoxyethyl mercapto)acetyl amino) adipoylglycyl]glycine. The  $^{186}\text{Re}$  complex formed is referred to as a  $^{186}\text{Re}$ -N<sub>3</sub>S complex, since



the  $^{186}\text{Re}$  is coordinated by the three nitrogens and the one sulfur atom of the ligand. The  $^{186}\text{Re}$  complex is formed by reducing 0.5 ml of  $^{186}\text{Re}$  perrhenate (400 mCi, 4 mCi/ $\mu\text{g}$  Re) contained in a purged 10 ml vial with 0.5 ml of a deoxygenated stock solution of 1 mg stannous chloride, 10 mg gentisic acid, and 250 mg citric acid. The solution is kept at room temperature for 15 minutes. To this solution is added 0.2 ml of a 1 mg ligand/1 ml isopropanol stock solution. The solution is heated for 15 minutes at 80°C.

5 A solution of 25 mg of protein in 1.0 ml PBS is added to the vial containing the  $^{186}\text{Re-N}_3\text{S}$  complex. The pH of the solution is adjusted to pH 9-9.5 with a 0.2 M sodium carbonate buffer. The solution is incubated at room temperature for 15 minutes. Given a typical conjugation yield of 50%, this reaction yields  $^{186}\text{Re-N}_3\text{S-NR-Lu-10}$  at a specific activity of 200 mCi  $^{186}\text{Re}/25$  mg NR-Lu-10.

10 The  $^{186}\text{Re-N}_3\text{S-NR-Lu-10}$  is purified using a PD-10 column (G-25M Sephadex PD-10 column; Pharmacia LKB Biotechnology Inc.; Piscataway, NJ) eluting with PBS, pH 7.4, collecting 0.5 ml fractions. The  $^{186}\text{Re-N}_3\text{S-NR-Lu-10}$  elutes in fractions 6 and 7, which are subsequently combined. Half of this solution is transferred to a vial containing 10 mg of gentisic acid.

20 The labeled protein is kept at room temperature and is monitored for the purity of  $^{186}\text{Re-N}_3\text{S-NR-Lu-10}$  using HPLC with a gel permeation column (Zorbax GF-250; 9.4 mm x 25.0 cm; DuPont, Wilmington, DE) eluting with 0.1 M phosphate, 0.1% sodium dodecyl sulfate, pH 6.7 mobile phase at a flow rate of 1 ml/min. Presence of autoradiolytic decomposition of the labeled monoclonal antibody is detected by the appearance of HPLC peaks that do not correspond to that for the  $^{186}\text{Re-N}_3\text{S-NR-Lu-10}$  monoclonal antibody. The radiolytic decomposition occurs in the solution that does not contain gentisic acid.

The invention may be embodied in other specific forms without departing from its spirit or essential characteristics. The described embodiments are to be considered in all respects only as illustrative and not restrictive. The scope of the invention is, therefore, indicated by the appended claims rather than by the foregoing description. All changes which come within the meaning and range of equivalency of the claims are to be embraced within their scope.

10       What is claimed is:

1. A composition for preparing a stable radiolabeled peptide comprising:

a peptide capable of being labeled with a radionuclide; and

5 a first stabilizer selected from the group consisting of gentisic acid, gentisyl alcohol and the water soluble salts, esters, derivatives, and mixtures thereof.

10 2. A composition as defined in claim 1, wherein the peptide is capable of being labeled with the radionuclide by use of a bifunctional chelate.

15 3. A composition as defined in claim 1, wherein the peptide is capable of being labeled with the radionuclide by direct labeling of the peptide.

4. A composition as defined in claim 1, wherein the peptide is capable of being labeled with the radionuclide  
20 by covalent binding of the radionuclide to the peptide.

5. A composition as defined in claim 1, further comprising a second stabilizer.

25 6. A composition as defined in claim 5, wherein the second stabilizer includes inositol.

7. A composition as defined in claim 5, wherein the second stabilizer includes ascorbic acid.

30

8. A composition as defined in claim 1, further comprising a buffer.

9. A composition as defined in claim 1, wherein the  
35 composition is in the form of a lyophilized kit.

10. A composition as defined in claim 1, wherein the peptide is labeled with a radionuclide.

11. A composition as defined in claim 10, wherein the  
5 composition is a liquid formulation.

12. A composition for preparing a stable radiolabeled protein comprising:

a protein capable of being labeled with a radionuclide; and

5 a first stabilizer selected from the group consisting of gentisic acid, gentisyl alcohol and the water soluble salts, esters, derivatives, and mixtures thereof.

10 13. A composition as defined in claim 12, wherein the protein is capable of being labeled with the radionuclide by use of a bifunctional chelate.

15 14. A composition as defined in claim 12, wherein the protein is capable of being labeled with the radionuclide by direct labeling of the peptide.

20 15. A composition as defined in claim 12, wherein the protein is capable of being labeled with the radionuclide by covalent binding of the radionuclide to the peptide.

16. A composition as defined in claim 12, further comprising a second stabilizer.

25 17. A composition as defined in claim 16, wherein the second stabilizer includes inositol.

30 18. A composition as defined in claim 16, wherein the second stabilizer includes ascorbic acid.

19. A composition as defined in claim 12, further comprising a buffer.

35 20. A composition as defined in claim 12, wherein the composition is in the form of a lyophilized kit.

21. A composition as defined in claim 12, wherein the protein is labeled with a radionuclide.

22. A composition as defined in claim 21, wherein the  
5 composition is a liquid formulation.

23. A composition for preparing a stable radiolabeled octreotide preparation comprising:

a quantity of DTPA-octreotide; and

5 a stabilizer selected from the group consisting of gentisic acid, gentisyl alcohol, and the water soluble salts, esters, derivatives, and mixtures thereof.

24. A composition as defined in claim 23, further  
10 comprising a second stabilizer.

25. A composition as defined in claim 24, wherein the second stabilizer includes inositol.

15 26. A composition as defined in claim 24, wherein the second stabilizer includes ascorbic acid.

27. A composition as defined in claim 24, further comprising a buffer.  
20

28. A composition as defined in claim 27, wherein the composition is in the form of a lyophilized kit.


29. A composition as defined in claim 27, wherein the  
25 DTPA-octreotide is labeled with a  $^{111}\text{In}$ .

30. A composition as defined in claim 29, wherein the composition is a liquid formulation.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 92/06429

<b>I. CLASSIFICATION OF SUBJECT MATTER</b> (If several classification symbols apply, indicate all) <sup>6</sup>		
According to International Patent Classification (IPC) or to both National Classification and IPC Int.Cl. 5 A61K49/02; A61K47/12		
<b>II. FIELDS SEARCHED</b>		
Minimum Documentation Searched <sup>7</sup>		
Classification System	Classification Symbols	
Int.Cl. 5	A61K	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched <sup>8</sup>		
<b>III. DOCUMENTS CONSIDERED TO BE RELEVANT<sup>9</sup></b>		
Category <sup>o</sup>	Citation of Document, <sup>11</sup> with indication, where appropriate, of the relevant passages <sup>12</sup>	Relevant to Claim No. <sup>13</sup>
Y	US,A,4 497 744 (FAWZI) 5 February 1985 cited in the application see column 3 - column 4; examples 1-10 ---	1-30
Y	EP,A,0 123 314 (NIHON MEDI-PHYSICS CO., LTD.) 31 October 1984 see example 3 ---	1-30
Y	US,A,4 232 000 (FAWZI) 4 November 1980 cited in the application see column 3 - column 4; claims 1-3; examples 1-10 --- -/--	1-30
<sup>o</sup> Special categories of cited documents : <sup>10</sup> "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. "&" document member of the same patent family		
<b>IV. CERTIFICATION</b>		
Date of the Actual Completion of the International Search	Date of Mailing of this International Search Report	
02 DECEMBER 1992	25. 01. 93	
International Searching Authority EUROPEAN PATENT OFFICE	Signature of Authorized Officer DULLAART A.W.M. 	



III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category °	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No.
Y	GB,A,2 239 178 (SANDOZ LTD) 26 June 1991 see page 13, paragraph 4 -paragraph 5 see page 14, paragraph 4 -paragraph 5 see page 18 - page 19 ---	1-30
Y	GB,A,2 225 579 (SANDOZ LTD) 6 June 1990 cited in the application see examples ---	1-30
P,X	WO,A,9 200 759 (MALLINCKRODT MEDICAL INC.) 23 January 1992 see page 12, line 20 - line 28; example 3 ---	1-28,30
P,X	LIFE SCIENCES, VOL. 49, NO. 22, PAGES 1593-1601, 1991, W.H. BAKKER ET AL. 'In vivo application of [111In-DTPA-D-Phe]-octreotide for detection of somatostatin-receptor positive tumors in rats.' see abstract ---	1-30
P,X	LIFE SCIENCES, VOL. 49, NO. 22, PAGES 1583-1591, 1991, W.H. BAKKER ET AL. '[111In-DTPA-D-Phe]-octreotide, a potential radiopharmaceutical for imaging of somatostatin-receptor positive tumors: synthesis, radiolabeling and in vitro validation.' see page 1587 see page 1589 ---	1-30
Y	EP,A,0 165 630 (MALLINCKRODT, INC) 27 December 1985 see abstract; claim 7 -----	1-30

**ANNEX TO THE INTERNATIONAL SEARCH REPORT**  
**ON INTERNATIONAL PATENT APPLICATION NO. US 9206429**  
**SA 63454**

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 02/12/92

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US-A-4497744	05-02-85	US-A- 4233284	11-11-80
		AU-B- 537689	05-07-84
		AU-A- 4559579	04-10-79
		CA-A- 1122524	27-04-82
		EP-A, B 0004684	17-10-79
		JP-C- 1383535	09-06-87
		JP-A- 55013258	30-01-80
		JP-B- 61048485	24-10-86
		JP-B- 1049329	24-10-89
		JP-C- 1572270	25-07-90
		JP-A- 61210042	18-09-86
EP-A-0123314	31-10-84	JP-C- 1609992	15-07-91
		JP-B- 2033692	30-07-90
		JP-A- 59199636	12-11-84
		AU-A- 2727384	01-11-84
		US-A- 4645660	24-02-87
US-A-4232000	04-11-80	AU-B- 521704	22-04-82
		AU-A- 4845979	03-01-80
		CA-A- 1118683	23-02-82
		EP-A, B 0006659	09-01-80
		JP-C- 1471137	14-12-88
		JP-A- 55036796	14-03-80
		JP-B- 63006528	10-02-88
GB-A-2239178	26-06-91	AU-B- 618270	19-12-91
		BE-A- 1001079	04-07-89
		CH-A- 677449	31-05-91
		DE-A- 3822557	19-01-89
		FR-A- 2617714	13-01-89
		GB-A, B 2208200	15-03-89
		JP-A- 1031728	02-02-89
		LU-A- 87268	08-03-89
		NL-A- 8801734	01-02-89
		SE-A- 8802569	11-01-89
GB-A-2225579	06-06-90	AU-A- 4587189	14-06-90
		BE-A- 1002296	20-11-90
		CA-A- 2004532	05-06-90

**ANNEX TO THE INTERNATIONAL SEARCH REPORT**  
**ON INTERNATIONAL PATENT APPLICATION NO. US 9206429**  
**SA 63454**

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report.  
The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information. 02/12/92

Page 2

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB-A-2225579		CH-A- 678329	30-08-91
		WO-A- 9006949	28-06-90
		FR-A- 2639947	08-06-90
		JP-A- 2184698	19-07-90
		NL-A- 8902981	02-07-90
		SE-A- 8904087	05-06-91
-----			
WO-A-9200759	23-01-92	AU-A- 8424291	04-02-92
-----			
EP-A-0165630	27-12-85	BE-A- 899739	26-11-84
		AU-B- 576861	08-09-88
		AU-A- 4272785	28-11-85
		US-A- 5021220	04-06-91
		US-A- 4942231	17-07-90
-----			